



Värdplantor för och skador av *H. schachtii* & *H. trifolii*

Crops	<i>H. schachtii</i> White BCN		<i>H. trifolii</i> Yellow BCN	
	Clay/ loamy/ sandy		Sandy soils	
Type of soil	multiplication	Crop damage	multiplication	Crop damage
Potato	-	not	-	not
Sugar beet	+++	strong	+++	strong
Onion	-	not	-	not
Maize	-	not	-	not
Winter wheat	-	not	-	not
Spring wheat	-	not	-	not
Winter barley	-	not	-	not
Spring barley	-	not	-	not
Rye	-	not	-	not
Oat	-	not	-	not
Triticale	-	not	-	not
Alfalfa	-	not	-	not
Flax	-	not	-	not
Winter canola	+++	a few	?	unknown
Summer canola	+++	strong	++	a few
Vegetable crops				
Pea (canning)	-	not	-R	strong
Dwarf French bean	-	not	+	average
Field-/gardenbean	-	not	++	average
Spinach	++	average	++	average
Carrot	-	not	-	not
Scorzonerä	-	not	-	not
Chicory	-	not	-	not
Cabbage species	+++	average	+++	average
Leek	-	not	-	not
Red beet	+++	strong	+++	strong
Flower bulbs				
Dahlia	-	not	-	not
Gladiolus	-	not	-	average
Lily	-	not	-	not
Tulip	-	not	-	not
Green manure crops				
Fodder radish	-	not	-	not
White mustard	-R	not	-R	not
Perennial ryegrass	-	not	-	not
Italian ryegrass	-	not	-	not
Rye	-	not	-	not
Clover	-	not	+	average
Lupine	?	unknown	?	unknown
Phacelia	-	not	-	not
Tagetes	-	not	-	not

Source: Nematode management in the agriculture. PPO-Lelystad 2000.

Legend:

Nematode multiplication	
?	Unknown
--	Active reduction
-	not
+	bad
++	average
+++	good
R	Depends on variety